

CVS Pharmacy, Inc.

-PE Product Labels –

VALUE PACK

NDC 69842-967-12

Compare to Mucinex® Fast-Max® Maximum Strength Severe Congestion & Cough Clear & Cool® Active Ingredients*

Maximum Strength Severe Congestion & Cough

Dextromethorphan HBr • Cough Suppressant
Guaifenesin • Expectorant
Phenylephrine HCl • Nasal Decongestant

- Controls Cough
- Relieves Nasal & Chest Congestion
- Thins & Loosens Mucus

Cooling Menthol Flavor

Naturality and Artificially Flavored

6 FL. OZ. (180 mL) For Ages 12+

Overnight Cold & Flu Clear & Cool

Acetaminophen • Pain Reliever/Fever Reducer
Dextromethorphan HBr • Cough Suppressant
Pseudoephedrine HCl • Nasal Decongestant
Tripropylene HD • Antihistamine

Maximum Strength per 4-hour dose

Night Time Relief for a Better Morning

- Cough • Fever • Sore Throat
- Runny Nose • Sneezing • Nasal Congestion

6 FL OZ (180 mL) For Ages 12+

MAXIMUM STRENGTH SEVERE CONGESTION & COUGH

OVERNIGHT COLD & FLU CLEAR & COOL

Active Ingredients	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant
Tripropylene HD 2 mg	Antihistamine

Drug Facts	Purposes
Active Ingredients (In each 20 mL) Dextromethorphan HBr 20 mg Guaifenesin 100 mg Phenylephrine HCl 10 mg Tripropylene HD 2 mg	Purposes Cough suppressant Expectorant Nasal decongestant Antihistamine

Uses

- Relieves throat irritation and soothes dry, scratchy throat
- Relieves chest and sinus congestion
- Relieves nasal congestion
- Relieves sore throat
- Relieves runny nose
- Relieves sneezing
- Relieves watery eyes
- Relieves itchy throat
- Relieves itchy nose
- Relieves itchy ears
- Relieves itchy skin
- Relieves itchy scalp
- Relieves itchy hair
- Relieves itchy nails
- Relieves itchy feet
- Relieves itchy hands
- Relieves itchy fingers
- Relieves itchy toes
- Relieves itchy palms
- Relieves itchy soles
- Relieves itchy heels
- Relieves itchy arches
- Relieves itchy insteps
- Relieves itchy backs
- Relieves itchy chests
- Relieves itchy abdomens
- Relieves itchy thighs
- Relieves itchy buttocks
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy shoulders
- Relieves itchy necks
- Relieves itchy faces
- Relieves itchy chins
- Relieves itchy cheeks
- Relieves itchy foreheads
- Relieves itchy brows
- Relieves itchy eyelids
- Relieves itchy eyebrows
- Relieves itchy hairlines
- Relieves itchy temples
- Relieves itchy ears
- Relieves itchy noses
- Relieves itchy mouths
- Relieves itchy lips
- Relieves itchy chins
- Relieves itchy necks
- Relieves itchy shoulders
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves itchy wrists
- Relieves itchy elbows
- Relieves itchy knees
- Relieves itchy hips
- Relieves itchy buttocks
- Relieves itchy thighs
- Relieves itchy legs
- Relieves itchy ankles
- Relieves it

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- Temporarily relieves:
 - minor aches and pains
 - headache
 - minor sore throat pain
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

- do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; stir while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 8 mg, sodium 20 mg
- phenylephrine HCl: contains phenylephrine 14 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

Inactive ingredients: Acesulfame K, Aspartame, Citric Acid, FD&C Blue No. 1, FD&C Red No. 40, Maltodextrin, Menthol, Natural Flavors, Sodium Citrate, Starch and Sugar.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2015 CVS/pharmacy
www.cvs.com 1-800-shop-CVS
Made in Canada
V-18533

CVS Quality
Money Back Guarantee

PLEASE RECYCLE

SEVERE COUGH & COLD DAYTIME

NDC 59779-220-06

CVS pharmacy

NEW!

SEVERE COUGH & COLD DAYTIME

ACETAMINOPHEN – PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr – COUGH SUPPRESSANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT

- Nasal & sinus congestion
- Cough
- Body ache
- Sore throat pain
- Headache
- Fever

Berry infused with Menthol & Green Tea flavors

6 PACKETS

SEE NEW WARNINGS INFORMATION

934160

0 50428 46341 5

C-3110-30508-6D

SEVERE COUGH & COLD DAYTIME

CVS/pharmacy™ Severe Cough & Cold Daytime provides complete relief of your cough and severe cold symptoms. This soothing hot liquid formula is alcohol free and tastes great.

For nighttime relief, try



Drug Facts (continued)

Directions

- do not take more than directed (see Overage warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cap provided
- do not use dosing cap with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cap provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 8 mg
- store at room temperature
- do not refrigerate

Inactive ingredients antemetic citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, glycerin, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose K100, polyethylene glycol 400, purified water, sodium benzoate, sorbitol, sucralose, sucrose, xanthan gum.

Questions or comments? 1-866-467-2748

2570

Drug Facts (continued)

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to 11 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose

Questions or comments?
Call 1-877-753-3936 Monday-Friday 9AM-5PM EST

CVS Health® Children's Cold & Allergy offers relief from stuffy nose, runny nose, sneezing, itchy nose and throat, and itchy, watery eyes to help your child get through the day.

Dosing Cup Included

***This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Allergy.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

IMPORTANT:
Keep this carton for future reference on full labeling.

Package Contains
One Bottle

Actual Size

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-13114

100% money back
guaranteed.
CVS.com/returnpolicy

Recycling symbols for Paper Box, Plastic Bottle, and Plastic Cup. Text: "Recycle Responsibly" and "Not recycled in all communities".

CVS Health.

Compare to the active ingredients in Children's Dimetapp® Cold & Allergy**

Grape Flavor

NDC 69842-779-04

Children's Cold & Allergy

BROMPHENIRAMINE MALEATE, 2 mg
Antihistamine
PHENYLEPHRINE HCl, 5 mg
Nasal decongestant

Liquid

Relieves:
Stuffy nose; Runny nose;
Sneezing; Itchy, watery eyes;
Itchy nose & throat

Alcohol free

Actual Bottle
Size on
Side Panel

Grape Flavor
For Ages 6 Years & Over

4 FL OZ (118 mL)

Drug Facts

Active Ingredients (in each 10 mL) **Purposes**
Brompheniramine maleate 2 mg Antihistamine
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis)
- runny nose ■ itchy, watery eyes
- sneezing ■ itching of the nose or throat
- temporarily restores free breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product ■ mL = milliliter


#477078

PLD-D2600 FC008254

Lot No.:

Exp. Date:





Compare to the active ingredients in Children's Mucinex® Multi-Symptom Cold*

NDC 59779-634-26

Children's Multi-Symptom Cold

DEXTROMETHORPHAN HBr 5 mg
Cough suppressant
GUAIFENESIN 100 mg
Expectorant
PHENYLEPHRINE HCl 2.5 mg
Nasal decongestant

Relieves:

- Stuffy nose
- Cough
- Chest congestion
- Breaks up mucus

Alcohol free



Very Berry Flavor Liquid
Ages 4+
Dosage cup included

4 FL OZ (118 mL)



Compare to the active ingredients in Children's Mucinex® Multi-Symptom Cold*

NDC 59779-634-26

Children's Multi-Symptom Cold

DEXTROMETHORPHAN HBr 5 mg
Cough suppressant
GUAIFENESIN 100 mg
Expectorant
PHENYLEPHRINE HCl 2.5 mg
Nasal decongestant

Relieves:

- Stuffy nose
- Cough
- Chest congestion
- Breaks up mucus

Alcohol free



Very Berry Flavor Liquid
Ages 4+
Dosage cup included

4 FL OZ (118 mL)

Multi-Symptom
Cold
Children's



Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 100 mg.....Expectorant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Drug Facts (continued)

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 3 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Multi-Symptom Cold.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

895141



LOT NO.

EXP.

Code Area

: 83926 17 C1





Compare to the active ingredients in Dimetapp® Cold & Cough*

Red Grape Flavor

Children's Cold & Cough

NDC 59779-363-26

BROMPHENIRAMINE MALEATE
Antihistamine
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant
DM Liquid

Relieves:
Nasal congestion; Stuffy nose;
Runny nose; Sneezes
Soothes cough
Calms allergy symptoms
Alcohol free



Red Grape Flavor
For Ages 6 Years & Over

4 FL OZ (118 mL)





Red Grape Flavor

CVS Health™ Children's Cold & Cough offers red grape-flavored relief from nasal congestion, runny nose, coughing and sneezing to get through the day.

Do not use if printed neckband is broken or missing

*CVS Health™ Children's Cold & Cough is not manufactured or distributed by Pfizer, distributor of Dimetapp® Cold & Cough.



PLEASE RECYCLE

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2016 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430



Money Back Guarantee



Children's Cold & Cough

DM Liquid

Red Grape Flavor

Drug Facts

Active Ingredients (In each 10 mL)
Brompheniramine maleate, USP 2 mg.....Antihistamine
Dextromethorphan HBr, USP 10 mg.....Cough suppressant
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Purposes
■ temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
■ temporarily relieves these symptoms due to hay fever (allergic rhinitis):
■ runny nose
■ sneezing
■ itchy, watery eyes
■ itching of the nose or throat
■ temporarily restores freer breathing through the nose

Uses
■ temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
■ temporarily relieves these symptoms due to hay fever (allergic rhinitis):
■ runny nose
■ sneezing
■ itchy, watery eyes
■ itching of the nose or throat
■ temporarily restores freer breathing through the nose

Warnings
Do not use
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ to make a child sleepy

Ask a doctor before use if you have
■ heart disease ■ high blood pressure
■ thyroid disease ■ diabetes
■ trouble urinating due to an enlarged prostate gland
■ glaucoma
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem such as emphysema or chronic bronchitis
■ cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
■ do not use more than directed
■ may cause marked drowsiness
■ avoid alcoholic beverages
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ excitability may occur, especially in children ▶

Drug Facts (continued)
Stop use and ask a doctor if
■ you get nervous, dizzy, or sleepless
■ symptoms do not get better within 7 days or are accompanied by fever
■ cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ do not take more than 6 doses in any 24-hour period
■ measure only with dosage cup provided
■ keep dosage cup with product
■ mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other Information
■ each 10 mL contains: sodium 4 mg
■ store at 20°-25°C (68°-77°F)

Inactive ingredients: anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?
1-800-719-9260

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

#477080-PE



0 50428 30748 9

LOT NO.

EXP.

: 94726 17 CL

No print/No varnish
Lot & Exp date

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each 5 mL)	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - stuffy nose
 - nasal congestion due to a cold
 - the intensity of coughing
 - the impulse to cough to help the child get to sleep

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has:

- diabetes
- heart disease
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor


Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C-red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-800-426-6361



TAMPER EVIDENT OR NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex® Children's Multi-Symptom Cold.

Package Contains One Bottle

Actual Size

CVS Health.

Children's Multi-Symptom Cold

DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
Oral Solution

VALUE SIZE

CVS Health.

Compare to the active ingredients in Mucinex® Children's Multi-Symptom Cold*

Children's Multi-Symptom Cold


DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
Oral Solution

Relieves:


- Chest congestion
- Stuffy nose
- Cough

Breaks up mucus
No added alcohol
Free of aspirin & ibuprofen

For Ages 4 to 11 Years



Berry Flavor
Dosage Cup Included



Actual Bottle Size on Side Panel

6.8 FL OZ (201 mL)

CVS Health.

Children's Multi-Symptom Cold

DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
Oral Solution

Relieves:


- Chest congestion
- Stuffy nose
- Cough

Breaks up mucus
No added alcohol
Free of aspirin & ibuprofen

For Ages 4 to 11 Years

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-19849

100% money back guaranteed.
CVS.com/returnpolicy



PAPER BOX PLASTIC BOTTLE PLASTIC CUP

B-0231-057-88-H
 REV0223B05788

#194705
FPO 100%
UPC# 050428600580
 X XXXXXX XXXXXX X

No Print/No Varnish
 Lot and Expiration No.



Compare to the active ingredients
in Mucinex® Fast-Max® Cold,
Flu & Sore Throat*

NO COPY AREA
Bottle curve zone

ADULT

Cold, Flu & Sore Throat

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MAXIMUM STRENGTH

- Relieves headache, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus

For Ages 12 & Over

6 FL OZ (180 mL)

NO COPY AREA
Bottle curve zone

: 16930 17 F1

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Do not use if printed neckband is broken or missing.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-16430

CVS Quality
Money Back Guarantee

NO COPY AREA
Bottle curve zone

#895139



0 50428 25876 7

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
 ■ cough ■ nasal congestion ■ minor aches and pains ■ headache
 ■ sore throat ■ temporarily reduces fever ■ temporarily promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

: 16930 17 F1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if ■ nervousness, dizziness or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

: 16930 17 F1

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 5 doses in any 24-hour period ■ measure only with dosing cup provided ■ do not use dosing cup with other products ■ dose as follows or as directed by a doctor ■ adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours ■ children under 12 years of age: do not use

Other information
 ■ each 20 mL contains: sodium 7 mg ■ dosing cup provided ■ store at 20-25°C (68-77°F) ■ do not refrigerate

Inactive ingredients anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triethyl citrate, xanthan gum

Questions or comments?
 1-800-719-9260

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Cold, Flu & Sore Throat.

: 16930 17 F1

Drug Facts	
Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 10 mg.....Cough suppressant	
Guaifenesin 200 mg.....Expectorant	
Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses	
■ temporarily relieves these common cold and flu symptoms	
■ headache ■ nasal congestion ■ sore throat	
■ cough ■ minor aches and pains	
■ temporarily reduces fever	
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	
Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
■ more than 4,000 mg of acetaminophen in 24 hours	
■ with other drugs containing acetaminophen	
■ 3 or more alcoholic drinks every day while using this product	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
■ skin reddening ■ blisters ■ rash	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use	
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
■ liver disease ■ diabetes ■ high blood pressure	
■ heart disease ■ thyroid disease	
■ trouble urinating due to an enlarged prostate gland	
Drug Facts (continued)	
■ a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
■ cough that occurs with too much phlegm (mucus)	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
When using this product, do not use more than directed.	
Stop use and ask a doctor if	
■ nervousness, dizziness, or sleeplessness occur	
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days	
■ fever gets worse or lasts more than 3 days	
■ redness or swelling is present	
■ new symptoms occur	
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
■ do not take more than directed (see Overdose warning)	
■ do not take more than 12 softgels in any 24-hour period	
■ adults and children 12 years of age and older: take 2 softgels every 4 hours	
■ children under 12 years of age: do not use	
Other information	
■ swallow whole; do not crush, chew, or dissolve	
■ avoid excessive heat	
■ store between 15-30°C (59-86°F)	
Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide	
Questions or comments?	
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	

CVS Health.

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

CVS Health.

Compare to the active ingredients in
Maximum Strength Mucinex® FAST-MAX®
Cold, Flu & Sore Throat*

NDC 69779-689-16

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

ACETAMINOPHEN - 325 mg - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - 10 mg - Cough suppressant
GUAIFENESIN - 200 mg - Expectorant
PHENYLEPHRINE HCl - 5 mg - Nasal decongestant

Relieves:

- Cough
- Nasal & chest congestion
- Headache & fever

Soothes cough
Thins & loosens mucusAges 12 Years & Over
Alcohol Free

Actual Size

16 LIQUID GELS
(Liquid filled capsules)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BILISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:

Exp. Date:

#177086



PLD-A42B FD03530

*This product is not manufactured or distributed by Robitussin. Because, distributor of Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
©2013 CVS Health. All rights reserved.
CVS Health® 1-800-Shop CVS V-16425CVS Quality
We care for your health.



Compare to the active ingredients
in Maximum Strength Mucinex®
FAST-MAX® Cold, Flu & Sore Throat*

No Print/No Varnish
Lot & Exp date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BULSTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING
B-0231-617-09-CFS
086091861709

MAXIMUM STRENGTH
Cold, Flu & Sore Throat
ACETAMINOPHEN
Pain reliever/Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant

Multi-Symptom

- Relieves aches, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus



Actual Size

20 CAPLETS

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever/fever reducer

Dextromethorphan HBr 10 mg.....Cough suppressant

Guaifenesin 200 mg.....Expectorant

Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ Temporarily relieves these common cold and flu symptoms:
■ nasal congestion ■ sore throat ■ headache ■ cough
■ minor aches and pains ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the
■ bronchial passages of bothersome mucus and make coughs more
productive ■ Temporarily reduces fever

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin reddening
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have
■ liver disease ■ diabetes ■ thyroid disease
■ heart disease ■ high blood pressure
■ difficulty in urination due to enlargement of the prostate gland

Drug Facts (continued)

Directions ■ do not take more than directed
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years and over: take 2 caplets every 4 hours
■ children under 12 years: do not use

Other information ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BULSTER IS TORN OR BROKEN
■ See end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients corn starch, croscarmellose, FD&C red #40, aluminum lake, FD&C yellow #6, aluminum lake, magnesium stearate, methocel K100, microcrystalline cellulose, polyethylene glycol, polyvinyl

Questions or comments? 1-800-426-6991

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-19849

CVS Quality
Money Back Guarantee

#322699

FPO 80%
UPC# 050428630426

X XXXXXXX XXXXXX X

CVS Health 44-617

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each 5 mL)	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - stuffy nose
 - nasal congestion due to a cold
 - the intensity of coughing
 - the impulse to cough to help the child get to sleep

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- diabetes
- heart disease
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C red #40, flavors, glycerin, polyethylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-800-426-9391

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Package Contains One Bottle

Actual Size

**Children's
Cough & Congestion**

DEXTROMETHORPHAN HBr
 Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl
 Nasal decongestant
Oral Solution

**Children's
Cough & Congestion**

DEXTROMETHORPHAN HBr
 Cough suppressant
GUAIFENESIN
 Expectorant
PHENYLEPHRINE HCl
 Nasal decongestant
Oral Solution

Relieves:

- Chest congestion
- Stuffy nose
- Cough

Breaks up mucus
 No added alcohol
 Free of high fructose corn syrup, sugar, aspirin & ibuprofen

For Ages 4 to 11 Years

Berry Flavor
 Dosage Cup Included
 Actual Bottle Size on Side Panel

6.8 FL OZ (201 mL)

**Children's
Cough & Congestion**

DEXTROMETHORPHAN HBr
 Cough suppressant
GUAIFENESIN
 Expectorant
PHENYLEPHRINE HCl
 Nasal decongestant
Oral Solution

Relieves:

- Chest congestion
- Stuffy nose
- Cough

Breaks up mucus
 No added alcohol
 Free of high fructose corn syrup, sugar, aspirin & ibuprofen

For Ages 4 to 11 Years

Distributed by:
 CVS Pharmacy, Inc.
 One CVS Drive
 Woonsocket, RI 02895
 © 2023 CVS/pharmacy
 CVS.com®
 1-800-SHOP CVS
 V-19849

100% money back guaranteed.
 CVS.com/returnpolicy

#194833

FPO 100%
UPC# 050428601785

X XXXXXX XXXXXX X

No Print/No Varnish
 Lot and Expiration No.

2581



Compare to the active ingredients
in Mucinex® FAST-MAX®
Congestion & Headache®

MAXIMUM STRENGTH

Congestion & Headache

ACETAMINOPHEN

Pain reliever/Fever reducer

GUAIFENESIN

Expectorant

PHENYLEPHRINE HCl

Nasal decongestant

Multi-Symptom

- Relieves headache & fever
- Relieves nasal & chest congestion
- Thins & loosens mucus



Actual Size

30 CAPLETS

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg Pain reliever/fever reducer
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)

Uses ■ Temporarily relieves these common cold and flu symptoms:
■ nasal congestion ■ fever ■ headache ■ minor aches and pains
■ sinus congestion and pressure
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the
bronchial passageways of bothersome mucus and make coughing more
productive ■ temporarily promotes nasal and/or sinus drainage

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage
may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms
may include: ■ blisters ■ rash ■ skin redness
If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription
or nonprescription). If you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)
(certain drugs for depression, psychiatric or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you
do not know if your prescription drug contains an MAOI, ask a doctor or
pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its
ingredients

Ask a doctor before use if you have ■ liver disease ■ diabetes
■ high blood pressure ■ heart disease ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic
bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)
**Ask a doctor or pharmacist before use if you are taking the blood thinning
drug warfarin.**

Drug Facts (continued)

Inactive ingredients corn starch, croscarmellose, FD&C red #40
aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate,
methylcellulose, microcrystalline cellulose, polyethylene glycol, polyvinyl
alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid,
talc, titanium dioxide

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No print/No varnish
Lot & Exp date

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com®
1-800-SHOP CVS V-19849



#927548

FPO 80%
UPC# 050428285107

X XXXXXX XXXXXX X

Drug Facts (continued)
Questions or comments? 1-800-426-4381

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex®
FAST-MAX® Congestion & Headache.
50844 ORG032061501

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed - see Overdose warning	
■ only use the dose cup provided	■ do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 41 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-9290

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe+VapoCOOL™.

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02896
© 2019 CVS/pharmacy
CVS.com® V-16430

CVS Health.

Compare to the active ingredients in Vicks® DayQuil® Severe+VapoCOOL™

NDC 69842-182-40

Cooling Severe Daytime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant

Relieves:

- Minor aches, pains & fever
- Nasal congestion & sinus pressure
- Cough
- Chest congestion

ALCOHOL 10%

12 FL OZ (355 mL)

Drug Facts DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

17 F2

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion and pressure
 - cough due to minor throat and bronchial irritation
 - minor aches and pains
 - headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depressive, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Other information

Each caplet contains sodium 3 mg.

Store at 20-25°C (68-77°F).

Use this product only if you have a sore throat, runny nose, cough, or cold. Do not use if you have a fever, sore throat, or cough that lasts more than 7 days, or if you have a rash or other allergic reaction to this product or any of its ingredients.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

Take only as directed - see overdose warning.

Do not exceed 8 caplets per 24 hrs.

Adults & children 12 yrs & over: 2 caplets with water every 4 hrs.

Children 6 to 11 yrs: 1 caplet with water every 4 hrs.

Children under 6 yrs: ask a doctor.

Store at 20-25°C (68-77°F).

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

CVS Health.

Compare to the active ingredients in Vicks® DayQuil® Severe+VapoCOOL™

NDC 69842-723-82

Cooling Severe Daytime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant

Relieves:

- Minor aches, pains & fever
- Nasal congestion & sinus pressure
- Cough
- Chest congestion

24 CAPLETS

Actual Size

PARENTS:

Learn about how this medicine works at www.StopMedicineAbuse.org

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe+VapoCOOL™.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Worcester, MA 02095
© 2019 CVS Pharmacy
CVS.com® 1-800-SHOP-CVS
V-66490

CVS Quality
Money Back Guarantee

#954405

50428 27109 4

35062 17 C1

Drug Facts (continued) **Warnings:** This product contains acetaminophen. Severe liver damage may occur if you take too much. Do not take more than 4,000 mg of acetaminophen in 24 hours. Do not drink alcohol while taking this product. **Directions:** See back panel for full directions. **Other information:** Contains 4 tablets. Each tablet contains 325 mg of acetaminophen. **Keep out of reach of children.** © 2019 CVS. All rights reserved. **CVS.com** **521000**

Drug Facts (continued)
Directions ■ Take orally as directed - see overdose warning
 ■ do not exceed 8 capsules per 24 hrs.
 adults & children 12 yrs. & over 2 capsules with water or beer, tea.

PARENTS:
Learn about teen medicine abuse
www.ShopMedicineAbuse.org

**DONOT USE IF PRINTED MEDICATION IS
BROKEN OR MISSING**
*These products are not manufactured or
distributed by Procter & Gamble, distributor
of Vicks® DayQuil® Severe+VapoCOOL™ and
Vicks® NyQuil® Severe+VapoCOOL™.

Drug Facts (continued)
Inactive ingredients: atchid, anhydrous citric acid, D&C yellow #10,
ediate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol,
propylene glycol, purified water, saccharin sodium, sodium benzoate,
sodium chloride, sodium citrate, sorbitol solution, sucralose
Questions or comments? 1-800-719-9280

CVS
Health™

Compare to the active
ingredients in Vicks® DayQuil®
Severe+VapoCOOL™*

CVS
Health™

Compare to the active
ingredients in Vicks® NyQuil®
Severe+VapoCOOL™**
NDC 69842-737-02

Cooling Severe Daytime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant

Relieves:

- Minor aches, pains & fever
- Nasal congestion & sinus pressure
- Cough
- Chest congestion

ALCOHOL 10%



12 FL OZ (355 mL)

Cooling Severe Nighttime

COLD & FLU RELIEF

ACETAMINOPHEN - Pain reliever/Fever reducer
PHENYLEPHRINE HCl - Nasal decongestant
DOXYLAMINE SUCCINATE - Antihistamine
DEXTROMETHORPHAN HBr - Cough suppressant

Relieves:

- Minor aches, pains & fever
- Nasal congestion & sinus pressure
- Sneezing, runny nose
- Cough

ALCOHOL 10%



12 FL OZ (355 mL)

Drug Facts (continued)

Inactive ingredients: alcohol, anhydrous citric acid, D&C yellow #10,
ediate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol,
propylene glycol, purified water, saccharin sodium, sodium benzoate,
sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-9280

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

CVS Quality
Money Back Guarantee

Nighttime		Daytime											
Drug Facts	Purpose	Drug Facts	Purpose										
Active ingredients (in each 30 mL)		Active ingredients (in each 30 mL)											
Acetaminophen 650 mg	Pain reliever/fever reducer	Acetaminophen 650 mg	Pain reliever/fever reducer										
Dextromethorphan HBr 20 mg	Cough suppressant	Dextromethorphan HBr 20 mg	Cough suppressant										
Doxylamine succinate 12.5 mg	Antihistamine	Guafenesin 400 mg	Expectorant										
Phenylephrine HCl 10 mg	Nasal decongestant	Phenylephrine HCl 10 mg	Nasal decongestant										
Uses		Uses											
Temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> nasal congestion sinus congestion and pressure minor aches and pains headache sore throat runny nose and sneezing cough to help you sleep cough due to minor throat and bronchial irritation reduces swelling in nasal passages fever promotes nasal and/or sinus drainage temporarily restores free breathing through the nose 	Temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> nasal congestion sinus congestion and pressure cough due to minor throat and bronchial irritation minor aches and pains headache fever reduces swelling in nasal passages temporarily restores free breathing through the nose sore throat helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive promotes nasal and/or sinus drainage 												
Warnings		Warnings											
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 	Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 												
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include <ul style="list-style-type: none"> skin redness hives rash If a skin reaction occurs, stop use and seek medical help right away.	Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include <ul style="list-style-type: none"> skin redness hives rash If a skin reaction occurs, stop use and seek medical help right away.												
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.												
Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients. 	Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients. 												
Ask a doctor before use if you have <ul style="list-style-type: none"> liver disease heart disease high blood pressure breathing problem such as emphysema or chronic bronchitis thyroid disease diabetes glaucoma cough that occurs with too much phlegm (mucus) trouble urinating due to an enlarged prostate gland persistent or chronic cough such as occurs with smoking, asthma, or emphysema a sodium-restricted diet 	Ask a doctor before use if you have <ul style="list-style-type: none"> liver disease heart disease thyroid disease high blood pressure trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema diabetes a sodium-restricted diet 												
Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> taking sedatives or tranquilizers taking the blood thinning drug warfarin 	Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none"> taking sedatives or tranquilizers taking the blood thinning drug warfarin 												
When using this product <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children be careful when driving a motor vehicle or operating machinery marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks 	When using this product <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children be careful when driving a motor vehicle or operating machinery marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks 												
Stop use and ask a doctor if <ul style="list-style-type: none"> pain, nasal congestion, or cough gets worse or lasts more than 7 days you get nervous, dizzy or sleepless fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. 	Stop use and ask a doctor if <ul style="list-style-type: none"> you get nervous, dizzy or sleepless pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. 												
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.												
Directions <ul style="list-style-type: none"> take only as directed - see Overdose warning only use the dose cup provided do not exceed 4 doses per 24 hrs <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>30 mL every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults & children 12 yrs & over	30 mL every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use	Directions <ul style="list-style-type: none"> take only as directed - see Overdose warning only use the dose cup provided do not exceed 4 doses per 24 hrs <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>30 mL every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults & children 12 yrs & over	30 mL every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL every 4 hrs												
children 4 to under 12 yrs	ask a doctor												
children under 4 yrs	do not use												
adults & children 12 yrs & over	30 mL every 4 hrs												
children 4 to under 12 yrs	ask a doctor												
children under 4 yrs	do not use												
Other information <ul style="list-style-type: none"> each 30 mL contains sodium 4 mg store at 20-25°C (68-77°F) 	Other information <ul style="list-style-type: none"> each 30 mL contains sodium 41 mg store at 20-25°C (68-77°F). Do not refrigerate. 												

E3 27 204922

72V62 17 C2



Compare to the
active ingredients
in Vicks® NyQuil®
Severe+VapoCOOL™*

NDC 69842-590-40

Cooling Severe Nighttime

ACETAMINOPHEN-

Pain reliever/Fever reducer

PHENYLEPHRINE HCl-

Nasal decongestant

DOXYLAMINE SUCCINATE-

Antihistamine

DEXTROMETHORPHAN HBr-

Cough suppressant

Relieves:

Minor aches & pains,
Fever, Nasal congestion,
Sinus pressure, Sneezing,
Runny nose, Cough

COLD & FLU

ALCOHOL 10%

12 FL OZ (355 mL)

: 59B40 17 F3

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

**DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING**

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

**100% money back
guaranteed.**
CVS.com/returnpolicy

Empty &
Replace Cap



PLASTIC
BOTTLE



PLASTIC
CUP

how2recycle.info

Drug Facts

Active ingredients (in each 30 mL)

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ runny nose and sneezing ■ sore throat ■ cough to help you sleep ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs,
stop use and seek medical
help right away.

**PEEL BACK AT
CORNER FOR MORE
INFORMATION**

: 59B40 17 B1

Drug Facts
(continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ heart disease
■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma
■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

Drug Facts
(continued)

When using this product ■ **do not use more than directed** ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 42 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions?

1-800-719-9260

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe+ VapoCOOL™.

ADHESIVE AREA
NO COATING • NO VARNISH • NO TYPE

ADHESIVE AREA
NO COATING • NO VARNISH • NO TYPE



2592

3.5625 in

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed
• only use the dose cup provided • do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 4 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

Other information • each 30 mL contains: sodium 81 mg • store at room temperature. Do not refrigerate.

Inactive ingredients citric acid, D&C Yellow No.10, disodium edetate, FD&C Blue No. 1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions or comments? 1-866-467-2748

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe + VapoCOOL®.

LOT: 72712CVSLR

UNVARNISHED
AREA

EXP:

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-12431
 **100% money back
guaranteed.**
CVS.com/returnpolicy

 **CVS
Health**

Compare to the active
ingredients in Vicks® DayQuil™
Severe + VapoCOOL™

Cooling Severe Daytime

ACETAMINOPHEN-
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr-
Cough suppressant
GUAIFENESIN-
Expectorant
PHENYLEPHRINE HCl-
Nasal decongestant

Relieves: Minor aches & pains, Fever,
Nasal congestion, Sinus pressure,
Cough, Chest congestion

COLD & FLU

12 FL OZ (354 mL)

Drug Facts

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SHRINK BAND IS MISSING OR BROKEN.**

Active ingredients (in each 30 mL)

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages
• temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hrs, which is the maximum daily amount for this product • taken with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.


Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • a sodium-restricted diet











2598

7.25 in

PDP 2.07 in

3.5625 in

Drug Facts (continued)

When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed

• only use the dose cup provided • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over 30 mL every 4 hrs

children 4 to under 12 yrs ask a doctor

children under 4 yrs do not use

Other information • each 30 mL contains: sodium 81 mg • store at room temperature. Do not refrigerate.

Inactive ingredients citric acid, D&C Yellow No.10, disodium edetate, FD&C Blue No.1, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose.

Questions or comments? 1-866-467-2748

Lot: 61412CVSLR

UNVARNISHED
AREA

EXP:

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil™ Severe+VapoCOOL™.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-12431

100% money back
guaranteed.
CVS.com/returnpolicy



Compare to the active
ingredients in Vicks® NyQuil™
Severe+VapoCOOL™**

Cooling Severe Nighttime

ACETAMINOPHEN-

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr-

Cough suppressant

DOXYLAMINE SUCCINATE-

Antihistamine

PHENYLEPHRINE HCl-

Nasal decongestant

Relieves: Minor aches & pains, Fever,
Nasal congestion, Sinus pressure,
Sneezing, Runny nose, Cough

COLD & FLU

12 FL OZ (354 mL)

Drug Facts

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SHRINK BAND IS MISSING OR BROKEN.**

Active ingredients (in each 30 mL)

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland • a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin.

Aspirin
SAC

Drug Facts

Active Ingredients (in each packet)

Acetaminophen 325 mg	Pain reliever/Fever reducer
Pheniramine maleate 20 mg	Antihistamine
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains • headache • minor sore throat pain
 - nasal congestion • temporarily reduces fever
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose • sneezing • itchy nose and throat
 - itchy, watery eyes

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 packets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use • with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

• with any other product containing acetaminophen (see Overdose Warning).

Ask a doctor before use if you have

• liver disease	• heart disease	• high blood pressure
• thyroid disease	• diabetes	• glaucoma
• a breathing problem such as emphysema, asthma, or chronic bronchitis		
• trouble urinating due to an enlarged prostate gland		
• a sodium-restricted diet		

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers • the blood thinning drug warfarin.

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.**

Overdose Warning: Taking more than the recommended dose can cause serious health problems, including serious liver damage. In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see Overdose Warning)
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor
- adults and children 12 years of age and over: dissolve contents of one packet in 8 oz. of hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other Information

- each packet contains: sodium 66 mg
- store at controlled room temperature 20°-25°C (68°-77°F)

Inactive ingredients

Citric Acid, D&C Yellow 10, Ethylcellulose, FD&C Red 40, Flavor, Microcrystalline Cellulose, Povidone, Silicon Dioxide, Sodium Citrate, Starch, Sugar and Vitamin C.

Questions or comments? 1-800-925-4733

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/Pharmacy
www.cvs.com 1-800-shop-CVS
Made in Canada
V-18633

CVS Quality
Money Back Guarantee

COLD & SORE THROAT
LEMON FLAVOR

NDC 59779-410-06

CVS pharmacy

NEW!

COLD & SORE THROAT
LEMON FLAVOR

PAIN RELIEVER / FEVER REDUCER (ACETAMINOPHEN)
ANTIHISTAMINE (PHENIRAMINE MALEATE)
NASAL DECONGESTANT (PHENYLEPHRINE HCL)

- Sore throat
- Nasal congestion
- Runny nose
- Headache
- Body ache
- Fever

6 PACKETS
Lemon Flavor

SEE NEW WARNINGS INFORMATION



CVS/pharmacy™ Cold and Sore Throat Relief provides temporary relief of cold symptoms, including sore throat, nasal congestion, runny nose, headache, body ache, and fever. This soothing hot liquid formula is alcohol free and tastes great.

834161



0 50428 46292 8

C-3173-30508

Nighttime Severe Cold & Cough

Drug Facts

Active ingredients (in each packet) **Purpose**
 Acetaminophen 650 mg **Pain reliever/fever reducer**
 Dextromethorphan hydrobromide 25 mg **Cough suppressant**
 Phenylephrine hydrochloride 10 mg **Nasal decongestant**

Uses

- Temporarily relieves:
 - minor aches and pains • headache • minor sore throat pain
 - nasal and sinus congestion • cough due to minor throat and bronchial irritation • runny nose • sneezing • itchy, watery eyes
 - due to hay fever • itchy nose and throat
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount, with other drugs containing acetaminophen, 3 or more alcoholic drinks every day while using this product. **See threat warning:** If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- if you are now taking a prescription non-narcotic cough suppressant (MAD) or have taken one in the last 14 days. If you do not know if your prescription drug contains a MAD, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are

- taking isotretinoin or tetracyclines • taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • rapid development may occur • drowsiness and dizziness may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children

Stop use and ask a doctor if

- nausea, vomiting, or sleepiness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worse • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Daytime Severe Cold & Cough

Drug Facts

Active ingredients (in each packet) **Purpose**
 Acetaminophen 650 mg **Pain reliever/fever reducer**
 Dextromethorphan hydrobromide 20 mg **Cough suppressant**
 Phenylephrine hydrochloride 10 mg **Nasal decongestant**

Uses

- Temporarily relieves:
 - minor aches and pains • minor sore throat pain
 - headache • nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount, with other drugs containing acetaminophen, 3 or more alcoholic drinks every day while using this product. **See threat warning:** If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- if you are now taking a prescription non-narcotic cough suppressant (MAD) or have taken one in the last 14 days. If you do not know if your prescription drug contains a MAD, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • rapid development may occur • drowsiness and dizziness may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children

Stop use and ask a doctor if

- nausea, vomiting, or sleepiness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worse • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Daytime

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Phenylpropanolamine is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not use more than directed
- Take every 4 hours, not to exceed 6 packets in 24 hours or as directed by a doctor
- Adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. of water, sip while hot. Consume entire drink within 15-15 minutes.
- Children under 12 years of age: consult a doctor
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briefly before and after heating. Do not overheat.

Other information

- Each packet contains: acetaminophen 650 mg, sodium 30 mg
- phenylephrine hydrochloride 10 mg
- phenylephrine hydrochloride 10 mg per packet
- Store at controlled room temperature 20°-25° (68°-77°F). Protect from moisture, heat, and moisture.

Inactive ingredients: Acetaminophen, acetaminophen, citric acid, FD&C Blue No. 1, FD&C Red No. 40, maltodextrin, mannitol, natural flavors, sodium citrate, starch and sugar.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Distributed by: CVS Pharmacy, Inc., One CVS Drive, Woonsocket, RI 02893
 © 2012 CVS-pharmacy
 www.cvs.com 1-800-450-CVS
 Made in Canada

CVS Quality
 Money Back Guarantee

CVS pharmacy

SEVERE COLD & COUGH DAYTIME

ACETAMINOPHEN - PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

Nasal & sinus congestion
Sore throat pain
Cough
Body ache
Headache
Fever

Berry infused with Mandarin & Green Tea Flavors

6 DAYTIME PACKETS

SEE NEW WARNINGS INFORMATION

CVS pharmacy

SEVERE COLD & COUGH NIGHTTIME

ACETAMINOPHEN - PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

Nasal congestion
Sore throat pain
Cough
Body ache
Headache
Fever
Runny nose
Sneezing

Honey Lemon infused with Chamomile & White Tea Flavors

6 NIGHTTIME PACKETS

12 PACKETS TOTAL

Nighttime

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Phenylpropanolamine is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not use more than directed
- Take every 4 hours, not to exceed 6 packets in 24 hours or as directed by a doctor
- Adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. of water, sip while hot. Consume entire drink within 15-15 minutes.
- Children under 12 years of age: consult a doctor
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briefly before and after heating. Do not overheat.

Other information

- Each packet contains: acetaminophen 650 mg, sodium 40 mg
- phenylephrine hydrochloride 10 mg per packet
- Store at controlled room temperature 20°-25° (68°-77°F). Protect from moisture, heat, and moisture.

Inactive ingredients: Acetaminophen, acetaminophen, citric acid, FD&C Blue No. 1, FD&C Red No. 40, maltodextrin, mannitol, natural flavors, sodium citrate, starch and sugar.


DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark "Theraflu".



FPD 100%
UPC# 050428391907





ADAPTIVE AREA

CVS Health. Compare to the active ingredients in Theraflu® ExpressMax® Daytime Severe Cold & Cough*

Severe Cold & Cough
ACETAMINOPHEN
 Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr
 Cough suppressant
PHENYLEPHRINE HCl
 Nasal decongestant

Relief of:
 Nasal congestion; Cough;
 Fever; Body ache;
 Sore throat; Headache

DAYTIME

Berry Flavor
 Alcohol content 10%
 For Ages 12 & Over

8.3 FL OZ (245 mL)

NDC 59779-691-08

PLD-C398B
 LB008884

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Daytime Severe Cold & Cough.

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2022 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS V-13114

#455446

0 50428 37797 0

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold
 - minor aches and pains
 - headache
 - nasal and sinus congestion
 - sore throat
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: sodium 16 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

Lot No.:
Exp. Date:

Daytime
Non-Drowsy
Multi-Symptom Cold/Flu Relief

ACETAMINOPHEN - Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
Aching & fever, sore throat, coughing,
and congested-stuffy head so you
can get through the day

Alcohol free
Antihistamine free




Actual Size

NDC 69842-742-16

Cold/Flu

Compare to the active ingredients in
Vicks® DayQuil® Cold & Flu LiquiCaps®†

CVS Health

†TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF ALUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts**Active ingredients (in each softgel)**

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Purposes**Drug Facts (continued)****Uses**

- temporarily relieves common cold and flu symptoms
- minor aches and pains ■ headache ■ sore throat
- nasal congestion ■ fever
- cough due to minor throat and bronchial irritation

Drug Facts (continued)**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ diabetes

- heart disease ■ thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Drug Facts (continued)**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients

butylated hydroxyanisole*, butylated hydroxytoluene*, FD&C red #40*, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide*, white ink

*contains one or more of these ingredients

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16425



PLEASE RECYCLE



PLD-F40R FC005131

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

#890425



0 50428 2587 1 2

CVS Health

Compare to the active ingredients in
Vicks® DayQuil® Cold & Flu LiquiCaps®

Cold/Flu

NDC 69842-848-24

Non-Drowsy

Daytime

Multi-Symptom Cold/Flu Relief

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
Aching & fever, sore throat, coughing,
and congested-stuffy head so you
can get through the day

Alcohol free
Antihistamine free



Actual Size

24 SOFTGELS** ("LIQUID-FILLED CAPSULES")

Drug Facts	
Active Ingredients (in each softgel)	Purposes
Acetaminophen 325 mg..... Pain reliever/fever reducer Dextromethorphan HBr 10 mg..... Cough suppressant Phenylephrine HCl 5 mg..... Nasal decongestant	
Uses	<ul style="list-style-type: none"> temporarily relieves common cold and flu symptoms: <ul style="list-style-type: none"> minor aches and pains headache sore throat nasal congestion fever cough due to minor throat and bronchial irritation
Warnings	<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. </p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters </p> <p>If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. </p> <p>Ask a doctor before use if you have <ul style="list-style-type: none"> liver disease diabetes heart disease thyroid disease high blood pressure trouble urinating due to an enlarged prostate gland persistent or chronic cough such as occurs with smoking, asthma, or emphysema cough that occurs with too much phlegm (mucus) </p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p>
Drug Facts (continued)	
<p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if <ul style="list-style-type: none"> pain, cough, or nasal congestion gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. </p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions <ul style="list-style-type: none"> do not take more than directed (see Overdose warning) do not take more than 4 doses in 24 hours adults and children 12 years and over: take 2 softgels with water every 4 hours swallow whole; do not crush, chew, or dissolve children under 12 years: do not use when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing </p> <p>Other information <ul style="list-style-type: none"> store between 15-30°C (59-86°F) avoid excessive heat </p> <p>Inactive ingredients butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, white ink</p> <p>Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p> <p><small>This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.</small></p>	

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS Pharmacy
CVS.com® 1-800-SHOP CVS
V-16425

CVS Quality
Money Back Guarantee

PLEASE RECYCLE

PLD-6969A F005699

0 50428 36348 5

#890395

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



3.5" height

Drug Facts (continued)

• new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • take only as directed • only use the dose cup provided • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over 30 mL every 4 hrs

children 6 to under 12 yrs 15 mL every 4 hrs

children 4 to under 6 yrs ask a doctor

children under 4 yrs do not use

Other information • each 15 mL contains: sodium 42 mg • store at room temperature. Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Yellow No. 6, flavor, glycerin, menthol, potassium citrate, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium chloride, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-866-476-2748

*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® DayQuil® Cold & Flu.

Distributed by CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-12431

100% money back guaranteed.
CVS.com/returnpolicy

Donated Recycled Plastic Cup
how2recycle.info

LOT: 61712CVSLR

UNVARNISHED
AREA

EXP:

VICKS
Health.

Compare to the active
ingredients in Vicks®
DayQuil® Cold & Flu*

Daytime

ACETAMINOPHEN-

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr-

Cough suppressant

PHENYLEPHRINE HCl-

Nasal decongestant

Relieves:

Headache, Fever, Sore throat, Minor aches & pains, Nasal congestion, Sinus pressure, Cough

COLD & FLU

12 FL OZ (354 mL)

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SHIRINK BAND IS MISSING OR BROKEN.

Active ingredients (in each 15 mL)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

• adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product • child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease • heart disease • high blood pressure

• thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema • a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if: you get nervous, dizzy or sleepless • pain, nasal congestion or cough get worse or lasts more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present

Do not print

2.07" PDP AREA

Drug Facts (continued)

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 2 mg
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, benzoic acid, citric acid, disodium EDTA, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium citrate

Questions or comments?

Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL
 AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2019 CVS/pharmacy
 CVS.com® 1-800-SHOP CVS V-13114



PLD-B319E LB005842



Daytime

NDC 69842-378-04

Children's Cold & Cough

DEXTROMETHORPHAN HBr

Cough suppressant

PHENYLEPHRINE HCl

Nasal decongestant

Relieves:

- Cough
- Stuffy nose

Alcohol 0.14%

For Ages

4 to 11 Years

Cherry Flavor



4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

Purposes

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use ■ in a child under 4 years of age ■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. **Ask a doctor before use if the child has** ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ cough that occurs with too much phlegm (mucus) ■ chronic cough that lasts, or as occurs with asthma. **When using this product, do not exceed recommended dosage. Stop use and ask a doctor if** ■ nervousness, dizziness or sleeplessness occurs ■ symptoms do not improve within 7 days or occur with fever ■ cough persists more than 7 days, comes back or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using

Nighttime Severe Cold (continued)

Drug Facts (continued)

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients DAC yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

Questions or comments?
Call 1-877-755-3635 Monday-Friday 9AM-5PM EST

Daytime Severe Cold

Drug Facts

Active ingredients (in each softgel)

Active ingredients	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
- headache ■ nasal congestion ■ sore throat
- cough ■ minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of toll-free mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness ■ hives ■ rash
- if a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease ■ diabetes ■ high blood pressure
- heart disease ■ thyroid disease
- trouble urinating due to an enlarged prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Daytime Severe Cold (continued)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if:

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitol, titanium dioxide

Questions or comments?
Call 1-877-755-3635 Monday-Friday 9AM-5PM EST

DAY & NIGHT COMBO PACK

CVS Health

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Day Severe Cold Liquid Gels and Night Cold & Flu Liquid Gels†
NDC 69842-906-24

MAXIMUM STRENGTH

Daytime Severe Cold

Acetaminophen
325 mg Pain reliever, Fever reducer

Dextromethorphan HBr
10 mg Cough suppressant

Guaifenesin
200 mg Expectorant

Phenylephrine HCl
5 mg Nasal decongestant

MULTI-SYMPTOM

- Relieves cough
- Thins & loosens mucus
- Relieves nasal & chest congestion
- Relieves aches, fever & sore throat

FOR AGES 12+ Actual Size

16 SOFTGELS™
24 TOTAL (16 Liqui-filled capsules)

MAXIMUM STRENGTH

Nighttime Severe Cold

Acetaminophen
325 mg Pain reliever, Fever reducer

Dextromethorphan HBr
10 mg Cough suppressant

Doxylamine Succinate
6.25 mg Antihistamine

Phenylephrine HCl
5 mg Nasal decongestant

MULTI-SYMPTOM

- Relieves cough
- Relieves runny nose & sneezing
- Relieves nasal congestion
- Relieves aches, fever & sore throat

FOR AGES 12+ Actual Size

8 SOFTGELS™

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-16425

CVS Quality
Money Back Guarantee

Lot No.:
Exp. Date:

#177184

0 50428 25842 2

PARENTS: FL0-D594A FC008258

Learn about how medicine works
www.CVSpharmacy.com/learn

Children's Cold & Cough Daytime

Drug Facts

Active Ingredients (in each 5 mL)
Dextromethorphan HBr 5 mg
Phenylephrine HCl 2.5 mg

Purposes
Cough suppressant
Nasal decongestant

Uses
temporarily relieve
nasal and sinus congestion
cough due to minor throat and bronchial irritation as may occur with a cold

Warnings
Do not use
in a child under 4 years of age
in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has
heart disease
high blood pressure
thyroid disease
diabetes
cough that occurs with too much phlegm (mucus)
chronic cough that lasts, or as occurs with asthma

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if
nervousness, dizziness, or sleeplessness occurs
symptoms do not improve within 7 days or occur with fever
cough persists more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
do not give more than 6 doses in any 24-hour period
measure only with dosing cup provided. Do not use any other dosing device.
keep dosing cup with product
mL = milliliter
shake well before using

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	5 mL every 4 hours
children under 4 years of age	do not use

Children's Cold & Cough Daytime (continued)

Drug Facts (continued)

Other Information
each 5 mL contains sodium 2 mg
store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acacia, ascorbic acid, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium citrate

Children's Nighttime Cold & Cough

Drug Facts

Active Ingredients (in each 5 mL)
Diphenhydramine HCl 1.25 mg
Phenylephrine HCl 2.5 mg

Purposes
Cough suppressant
Nasal decongestant

Uses
temporarily relieve
runny nose
itchy throat
itchy, watery eyes due to hay fever
nasal and sinus congestion
cough due to minor throat and bronchial irritation as may occur with a cold

Warnings
Do not use
in a child under 4 years of age
in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
with any other product containing diphenhydramine, even one used or sold
for the purpose of making your child sleepy

Ask a doctor before use if the child has
heart disease
high blood pressure
thyroid disease
diabetes
glaucoma
cough that occurs with too much phlegm (mucus)
chronic cough that lasts, or as occurs with asthma
a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

Children's Nighttime Cold & Cough (continued)

Drug Facts (continued)

When using this product
do not exceed recommended dosage
marked drowsiness may occur
excitability may occur, especially in children
sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if
nervousness, dizziness, or sleeplessness occurs
symptoms do not improve within 7 days or occur with a fever
cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
do not give more than 6 doses in 24 hours unless directed by a doctor
measure only with dosing cup provided. Do not use any other dosing device.
keep dosing cup with product
mL = milliliter
shake well before using

Age	Dose
children 6 to under 12 years of age	10 mL every 4 hours
children 4 to under 6 years of age	do not use unless directed by a doctor
children under 4 years of age	do not use

Other Information
each 5 mL contains sodium 3 mg
store between 20-25°C (68-77°F). Do not refrigerate.
protect from light

Inactive ingredients acacia, ascorbic acid, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?
Call 1-877-763-3635 Monday-Friday 9AM-5PM EST

PARENTS:
Learn about how to keep your child safe.
www.SuppliesHouse.org

DAY & NIGHT COMBO PACK

CVS Health.
Compare to the active ingredients in Children's Triaminic® Day Time Cold & Cough*

Daytime

Children's Cold & Cough
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Cough
Stuffy nose
Alcohol 0.14%
For Ages 4 to 11 Years



Cherry Flavor
Dosage cup included
Actual Bottle Size on Side Panel

CVS Health.
Compare to the active ingredients in Children's Triaminic® Night Time Cold & Cough*

Nighttime

Children's Cold & Cough
DIPHENHYDRAMINE HCl
Antihistamine; Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Cough
Runny, stuffy nose
Itchy throat
Alcohol free
For Ages 6 to 11 Years



Grape Flavor
Dosage cup included
Actual Bottle Size on Side Panel

Package Contains Two Bottles
Actual Size

**4 FL OZ (118 mL) + 4 FL OZ (118 mL)
TOTAL 8 FL OZ (236 mL)**

CVS Quality
Money Back Guarantee

**PLD-8319E
PLD-8320E
7/20/2006**

Lot No.:
Exp. Date:

#194300

0 50428 60148 8

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2019 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**



Nighttime Cold & Flu**Drug Facts****Active ingredients (in each 30 mL)**

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains
- fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed — see Overdose warning
- only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9260

Daytime Cold & Flu**Drug Facts****Active ingredients (in each 15 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion ■ cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains ■ fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed — see Overdose warning
- only use the dose cup provided ■ do not exceed 4 doses per 24 hrs


adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 7 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9260



Daytime Multi-Symptom

Relieves:
Aches, Fever, Cough;
Nasal congestion;
Nasal decongestant
PHENYLEPHRINE HCl
Cough suppressant
DEXTROMETHORPHAN HBr
Pain reliever, Fever reducer
ACETAMINOPHEN

COLD & FLU RELIEF

Daytime Multi-Symptom

Non-Drowsy

Nighttime Multi-Symptom

Relieves:
Aches, Fever, Cough;
Sore throat, Sneezing;
Runny nose
Antihistamine
DOXYLAMINE SUCCINATE
Cough suppressant
DEXTROMETHORPHAN HBr
Pain reliever, Fever reducer
ACETAMINOPHEN

COLD & FLU RELIEF

Nighttime Multi-Symptom

DAY & NIGHT TWIN PACK

CVS Health

Compare to the active ingredients
in Vicks® DayQuil® and NyQuil® LiquiCaps®

NDC 59779-646-24

CVS Pharmacy, Inc.
Pharmacy
300 SHOP CVS
#278531

0 5042855409 8

Nighttime Cold & Flu Softgel	Daytime Cold & Flu Softgel												
<p>Drug Facts</p> <p>Active ingredients (in each softgel)</p> <table style="width: 100%;"> <tr> <td style="width: 60%;">Acetaminophen 325 mg.....</td> <td style="width: 40%;">Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan HBr 15 mg.....</td> <td>Cough suppressant</td> </tr> <tr> <td>Doxylamine succinate 6.25 mg.....</td> <td>Antihistamine</td> </tr> </table> <p>Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	Acetaminophen 325 mg.....	Pain reliever/fever reducer	Dextromethorphan HBr 15 mg.....	Cough suppressant	Doxylamine succinate 6.25 mg.....	Antihistamine	<p>Drug Facts</p> <p>Active ingredients (in each softgel)</p> <table style="width: 100%;"> <tr> <td style="width: 60%;">Acetaminophen 325 mg.....</td> <td style="width: 40%;">Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan HBr 10 mg.....</td> <td>Cough suppressant</td> </tr> <tr> <td>Phenylephrine HCl 5 mg.....</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ headache ■ minor aches and pains ■ fever ■ sore throat</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	Acetaminophen 325 mg.....	Pain reliever/fever reducer	Dextromethorphan HBr 10 mg.....	Cough suppressant	Phenylephrine HCl 5 mg.....	Nasal decongestant
Acetaminophen 325 mg.....	Pain reliever/fever reducer												
Dextromethorphan HBr 15 mg.....	Cough suppressant												
Doxylamine succinate 6.25 mg.....	Antihistamine												
Acetaminophen 325 mg.....	Pain reliever/fever reducer												
Dextromethorphan HBr 10 mg.....	Cough suppressant												
Phenylephrine HCl 5 mg.....	Nasal decongestant												
<p>Drug Facts (continued)</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers</p> <p>When using this product ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery</p> <p>Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p>Drug Facts (continued)</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>												

<p>Antihistamine free</p> <p>16 LIQUID GELS</p> <p>Actual Size</p>	<p>8 LIQUID GELS 24 TOTAL (Liquid filled capsules)</p> <p>Actual Size</p>	<p>When using DayQuil products, careful to ensure correct use. PLD-A42R FC PARENTS: Learn about your medicine www.StopNasalSneezing.com Distributed by One CVS Drive © 2016 CVS/pharm CVS.com® 1-800-441-1010 V-16425</p>
<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none">■ do not take more than directed (see Overdose warning)■ do not take more than 4 doses in 24 hours■ swallow whole; do not crush, chew, or dissolve■ adults and children 12 years and over: take 2 softgels with water every 4 hours.■ children under 12 years: do not use <p>Other information</p> <ul style="list-style-type: none">■ store between 15-30°C (59-86°F) ■ avoid excessive heat <p>Inactive ingredients butylated hydroxyanisole*, butylated hydroxytoluene*, carminic acid*, D&C yellow #10*, edible white ink, FD&C red #40*, FD&C yellow #6*, gelatin, glycerin, polyethylene glycol*, povidone, propylene glycol, purified water, sodium metabisulfite*, sorbitan*, sorbitol</p> <p>Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none">■ do not take more than directed (see Overdose warning)■ do not take more than 4 doses in 24 hours■ swallow whole; do not crush, chew, or dissolve■ adults and children 12 years and over: take 2 softgels with water every 6 hours■ children under 12 years: do not use <p>Other information</p> <ul style="list-style-type: none">■ store between 15-30°C (59-86°F) ■ avoid excessive heat <p>Inactive ingredients D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan*, sorbitol *may contain this ingredient</p> <p>Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p> <p>This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter and Gamble Company.</p>	

When using daytime and nighttime softgels, carefully read the labeling to ensure correct dosing.

Actual Size

8 SOFTGELS**

24 TOTAL

(Liquid-filled capsules)**

Relieves:
Aches; Fever; Cough;
Sore throat; Sneezing;
Furry nose
Alcohol free

DOXYLAMINE SUCCINATE
Cough suppressant
Artihistamine

DEXTROMETHORPHAN HBr
Pain reliever; Fever reducer

ACETAMINOPHEN
COLD & FLU RELIEF

Nighttime

Multi-Symptom

NDC 69842-744-24

PARENTS:
Learn about your medicine online
www.StayMedicineAlive.org

PLD-9427 FD005121

KEEP OUT OF REACH OF CHILDREN. THIS PRODUCT CONTAINS A SCHEDULE II CONTROLLED SUBSTANCE.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16425

CVS Quality
Money Back Guarantee

#164257

0 50428 55720 4

of The Procter and Gamble Company.

Actual Size

16 SOFTGELS**

Relieves:
Aches; Fever; Sore throat;
Cough; Nasal congestion
Alcohol free
Antihistamine free

DEXTROMETHORPHAN HBr
Pain reliever; Fever reducer

ACETAMINOPHEN
COLD & FLU RELIEF

Daytime

Multi-Symptom

Non-Drowsy

CVS Health™

DAY & NIGHT TWIN PACK

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®†

Nighttime Cold & Flu Softgel		Daytime Cold & Flu Softgel	
Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 15 mg.....Cough suppressant Doxylamine succinate 6.25 mg.....Antihistamine		Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant	
Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing		Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ headache ■ minor aches and pains ■ fever ■ sore throat	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Drug Facts (continued) Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.		Drug Facts (continued) Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland		Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema	
Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers		Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
When using this product ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery		When using this product, do not use more than directed.	
Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.		Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	

Daytime Cold & Flu Softgel	
Drug Facts (continued)	
Directions ■ do not take more than 4 doses in 24 hours ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ children under 12 years: do not use	
Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat	
Inactive ingredients butylated hydroxyanisole*, butylated hydroxytoluene*, FD&C red #40*, FD&C yellow #6*, gelatin, glycerin, polyethylene glycol*, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide*, white ink *may contain this ingredient	
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	
This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquidCap® are registered trademarks	

Nighttime Cold & Flu Softgel	
Drug Facts (continued)	
Directions ■ do not take more than 4 doses in 24 hours ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use	
Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat	
Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide*, white ink *may contain this ingredient	
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	
This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquidCap® are registered trademarks	